

Archean to Proterozoic depletion in Cape Verde lithospheric mantle

C. BONADIMAN,¹ M. COLTORTI¹, F. SIENA¹,
S. Y. O'REILLY², W. L. GRIFFIN² AND N. J. PEARSON²

¹ Department of Earth Sciences, University of Ferrara, 44100 Ferrara, Italy; costanza.bonadiman@unife.it

² GEMOC Key Centre, Dept. Earth and Planetary Sciences, Macquarie University, Sydney, NSW 2109, Australia

Two groups of protogranular spinel-lherzolite and harzburgite xenoliths occur in Late Tertiary necks on Sal Island, Cape Verde Archipelago. Large protogranular clinopyroxenes from lherzolites (cpx 8-17vol%) have high Cr₂O₃ and low CaO contents, and upward-convex Rare Earth Element (REE) patterns (LaN=3.6-5.3; SmN= 8.3-12.8; YbN=2.4-4.8). Orthopyroxenes show very high Cr₂O₃ contents and flat HREE distributions, with YbN ranging from 0.7 to 0.9. Small secondary clinopyroxenes in harzburgites (cpx 1-3vol%) are characterized by strong enrichment of LREE and MREE (LaN=7.9-64.1; SmN=36.6-95.4) with positively fractionated HREE patterns (YbN=4.9-13.1).

Orthopyroxenes in harzburgites reflect high degrees of partial melting, with low Al₂O₃ contents (<2.5 wt%), and negatively fractionated HREE patterns (YbN=0.7-0.9; SmN=0.07-0.13). Whole-rock and mineral chemistry strongly support two different origins for the two lithotypes, which cannot be ascribed to a simple progressive depletion process. The Cape Verde harzburgites were produced by high -degree partial melting (>25%) of a spinel-bearing protolith, whereas the lherzolites reflect low-degree partial melting of a garnet-bearing lherzolite that re-equilibrated in the spinel stability field.

Both lithotypes have been metasomatised by kimberlite-like melts, leaving veins of K-rich glass+K-feldspar. In situ Re/Os analyses of intergranular sulfides from lherzolites yield values of ¹⁸⁷Os/¹⁸⁸Os varying from 0.1037 ± 0.0020 to 0.1256 ± 0.0022. TRD model ages define three groups: 3320 ± 150 Ma, 2060 ± 140 Ma and 996 ± 73 Ma (1σ).

The age patterns in the Re-Os data, the juxtaposition of strongly depleted harzburgites and (originally) garnet lherzolites, and evidence of kimberlite-like metasomatism suggest that part of the ancient African subcontinental lithospheric mantle was incorporated into the newly formed oceanic lithosphere during the opening of the Atlantic ocean.