

Evolution of the Vestmannaeyjar volcanic system

I.A. SIGURDSSON¹ AND S.P. JAKOBSSON²

¹ South Iceland Nature Centre, ingvar@nattsud.is

² Icelandic Institute of Natural History, sjak@ni.is

The Vestmannaeyjar archipelago forms the southernmost tip of the Eastern Volcanic Zone in Iceland. The volcanic islands (largest are Heimaey, 13.4 km² and Surtsey, 1.4 km²), along with numerous submarine eruption sites, constitute the Vestmannaeyjar volcanic system. It has been suggested that the age of the Vestmannaeyjar system is about 0.1 Ma and that the Heimaey area is evolving into a central volcano [1].

A detailed mapping of Heimaey indicates that the island is built up of 12 eruptive units, ranging in composition from alkali olivine basalt to mugearite. We have sampled the island with emphasis on the complex Nordurklettur Formation of N-Heimaey and so far 31 rock samples have been analyzed for major and trace elements.

The Vestmannaeyjar rock suite divides into three groups based on mineralogy and rock chemistry. These are two groups of alkali basalts, VE I with MgO 8-12 wt% and VE II with MgO 5.5-7.7 wt%; and a group of hawaiites-mugearites, VE III with MgO 3.2-5.2 wt% [1]. The VE II basalts can be divided further into two groups based on trace elements where the older basalts are slightly more enriched in the more incompatible trace elements at the same MgO content.

There are strong indications that the volcanism on Heimaey has been cyclic, where each cycle starts with a VE I basalt, followed by the more evolved VE II basalt and ending with a hawaiite or mugearite VE III. During the Holocene one such cycle starts with the Storhöfði eruption about 6500 BP, followed by the Saeðjall and Helgafell eruptions approximately 6200 BP and ending with the Eldfell eruption in AD 1973. In the Nordurklettur Formation there appears to be at least one other comparative cycle present.

Four samples from the Nordurklettur Formation are currently being dated by the Ar-Ar method in order to get a better understanding of the inferred cyclic activity. Three drillcores from a 1600 m deep hole on Heimaey are also being dated; the samples are from depths of 905, 1010 and 1517 m [2]. In 2005 a new 2277 m deep hole was drilled on the eastern part of Heimaey [3]. Drill cuttings from this hole are being analyzed for major elements and will be analyzed for trace elements later in the year. In this and subsequent study we will try to determine more precisely the onset and evolution of the Vestmannaeyjar volcanic system.

References

- [1] Jakobsson S.P., (1979) *Acta Nat. Isl.* **26**, 106 p.
- [2] Palmason G. et al. (1965) *Nat. Energy Auth.*, 43 p.
- [3] Franzson H., (2006, in prep.).