

The halogen cycle in subduction zones: Insight from back-arc basin basalts

D. CHAVRIT^{1*}, L. RUZIE¹, R. BURGESS¹, D.R. HILTON²,
H. SUMINO³, J. SINTON⁴ AND C.J. BALLENTINE¹

¹S.E.A.E.S., University of Manchester, Manchester, UK

(*presenting author, deborah.chavrit@manchester.ac.uk)

²Scripps Institution of Oceanography, La Jolla, USA

³GCRC, University of Tokyo, Tokyo 113-0033, Japan

⁴University of Hawaii, Honolulu, USA

The extent to which the subduction process preserves the volatile elements signature of the downgoing slab and the mechanisms by which these elements are transferred into the mantle wedge are not well understood. Halogens (Cl, Br, I) are good candidates to trace these processes, due to their incompatibility and their relatively high concentrations in seawater and marine sediments. A technique developed at the University of Manchester [1] allows the high precision measurements of these elements on neutron-irradiated samples using noble gas mass spectrometry.

To better constrain the cycle of halogens in subduction zones, we analyzed the halogens in 15 volcanic glasses (BABB) from three back-arc basins which are known to contain slab-derived components *viz* Manus basin, Lau basin and Mariana trough.

The three back-arc basins have relatively constant Br/Cl weight ratios ($4.0 \pm 0.4 \times 10^{-3}$) which are 2x higher than the mid-ocean ridge basalts (MORB) value [2]. The I/Cl weight ratios (0.9 to 7.1×10^{-5}) range from values close to seawater to MORB values. These results suggest that the halogen composition of the BABB mantle source is affected by a slab-derived component. However, the I/Cl ratios positively correlate with Ba/Nb ratios that are between 5-33 (weight), which reflect the extent of the slab contribution. Thus, it indicates the presence of an unknown end member with a MORB-like Ba/Nb ratio and with low I/Cl and high Br/Cl ratios. It is notable that the halogen ratios of this component are similar to that of the fluid phases trapped in altered oceanic crust [3]. Another component with higher Ba/Nb, higher I/Cl and lower Br/Cl ratios, is consistent with the presence of a sedimentary-derived component. The possible origins of the signature of the halogen BABB mantle source will be discussed by comparing with the different components characterizing the subducted oceanic crust.

[1] Johnson *et al.* (2000), GCA **64**, 717-732; [2] Ruzié *et al.* (2012), V31A-2762 AGU Fall Meeting; [3] Chavrit *et al.* (2012), Mineral. Mag. **76** (6), 1566.

A new approach for LA-ICP-MS using a high sensitive mass spectrometer

R. CHEMNITZER^{1*} AND M. HAMESTER¹

¹Bruker Daltonik GmbH, Fahrenheitstr. 4, 28359 Bremen, Germany

(*correspondence: Rene.Chemnitzer@bruker.com),
(Meike.Hamester@bdal.de)

The geochemical characterization of samples includes the elemental distribution but also the isotopic composition for a broad range of elements. Laserablation coupled to an ICP-MS has become an indispensable method. New instrumental developments allow resolutions to single-digit μm spots and new ablation cells show improved transport characteristics.

With a high sensitive ICP-MS (Bruker auroraElite) these developments lead to a tremendous increase in information. The work presents results from a LA-ICP-MS setup that provides the highest sensitivity currently available and shows results of the quantitative and isotope ratio analysis of zircon samples and other minerals.

Different parameters were investigated:

Laser

- Spot size (Laser)
- Energy density

ICP-MS

- Sample introduction / plasma
- 3-D ion focusing
- Scan speed

The results verify that the considerable higher sensitivity leads to new alternative approaches in LA-ICP-MS, that will be discussed in detail.