

## **Impact-driven ocean acidification as a mechanism of Cretaceous–Palaeogene mass extinctions**

SOHSUKE OHNO<sup>1</sup>, TOSHIHIKO KADONO<sup>2</sup>,  
KOSUKE KUROSAWA<sup>1</sup>, TAIGA HAMURA<sup>3</sup>,  
TATSUHIRO SAKAIYA<sup>4</sup>, KEISUKE SHIGEMORI<sup>5</sup>,  
YOICHIRO HIRONAKA<sup>5</sup>, TAKAYOSHI SANNO<sup>5</sup>,  
TAKESHI WATARI<sup>5</sup>, KAZUTO OTANI<sup>6</sup>,  
TAKAFUMI MATSUI<sup>1</sup> AND SEIJI SUGITA<sup>3</sup>

<sup>1</sup>Planetary Exploration Research Center, Chiba Institute of Technology, Chiba 275-0016, Japan

<sup>2</sup>University of Occupational and Environmental Health, Kitakyushu 807-8555, Japan

<sup>3</sup>Department of Complexity Science and Engineering, University of Tokyo, Kashiwa 277-8561, Japan

<sup>4</sup>Department of Earth and Space Science, Graduate School of Science, Osaka University, Osaka 560-0043, Japan

<sup>5</sup>Institute of Laser Engineering, Osaka University, Osaka 565-0871, Japan

<sup>6</sup>Institut national de la recherche scientifique – Énergie Matériaux Télécommunications, Varennes J3X 1S2 Canada

The Cretaceous–Paleogene (K–Pg) mass extinction event at 65.5 Ma triggered by a meteorite impact is one of the most drastic events in the history of life on the Earth. Many hypotheses have been proposed as killing mechanisms induced by the impact, including global darkness due to high concentrations of atmospheric silicate dust particles, global wildfires, greenhouse warming due to CO<sub>2</sub> release, and global acid rain. However, the actual mechanism of extinction remains highly controversial. One of the most important clues for understanding the extinction mechanism is the marine plankton record, which indicates that plankton foraminifera, living in the near-surface ocean, suffered very severe extinction in contrast to the high survival ratio of benthic foraminifera. No proposed extinction mechanism can account for this globally observed marine extinction pattern. Here, we show that SO<sub>3</sub>-rich impact vapor was released in the K–Pg impact and resulted in the occurrence of global acid rain and sudden severe ocean acidification at the end of the Cretaceous, based on the new results of impact experiments at velocities much higher than previous works (> 10 km/s) and theoretical calculations on aerosol coagulation processes. Sudden severe ocean acidification can account for many of the features of various geologic records at the K–Pg boundary, including severe extinction of plankton foraminifera.