

Dissolution of K-feldspar at CO₂-saturated conditions

JÖRGEN ROSENQVIST^{1*}, ANDREW D. KILPATRICK²,
BRUCE W.D. YARDLEY²
AND CHRISTOPHER A. ROCHELLE¹

¹British Geological Survey, Keyworth, Nottingham NG12
5GG, UK (*correspondence: jorenq@bgs.ac.uk,
caro@bgs.ac.uk)

²School of Earth and Environment, University of Leeds, Leeds
LS2 9JT, UK
(ee07adk@leeds.ac.uk, b.w.d.yardley@leeds.ac.uk)

Aquifers in deep sedimentary basins have been identified as suitable targets for geological carbon dioxide storage, especially aquifers located in sandstone host rock. This has led to renewed interest in studying the interaction between sandstone minerals and aqueous fluids, as there is a paucity of data for CO₂-containing systems at relevant conditions. In an attempt to improve data coverage for important silicate minerals, we have measured the dissolution kinetics of K-feldspar in CO₂-saturated fluids over a range of temperatures.

K-feldspar grains were reacted with water in batch-type reactors at temperatures from 20°C to 200°C and pressures up to 200 bar. The mineral grains were allowed to react with pure water for a number of weeks before injection of CO₂ into the system. Excess CO₂ was provided to ensure CO₂ saturation. While the reaction time before injection was not long enough to attain complete equilibrium, it did considerably lower the degree of undersaturation with respect to K-feldspar and helped highlight the effect of CO₂ injection into a rock-equilibrated aqueous fluid.

At all temperatures studied, injection of CO₂ resulted in a rapid increase in the soluble concentrations of K and Si (and also Na from a plagioclase component). The dissolution then reached apparent steady state conditions after a few days, with observed dissolution rates in the range of 10⁻⁹ to 10⁻¹² mol m⁻² s⁻¹ over the temperature range studied. The CO₂-saturated solutions maintained mildly acidic conditions throughout the experiments and the observed rates therefore fall roughly between rates measured in strong acids and rates measured in unbuffered aqueous solution. However, our rates are slightly slower than those predicted based solely on pH, a fact that we preliminary attribute to our choice to pre-react the mineral and fluid before the CO₂ injection. The exact cause of this effect is still under investigation.