

REE abundances in Ediacaran to Early Cambrian phosphate deposits, Yangtze Platform (South China) by LA-ICPMS

DOROTHEE HIPPLER^{1,4}, ANDREAS KLÜGEL²,
QINGJUN GUO³ AND GERHARD FRANZ⁴

¹Graz University of Technology, Institute of Applied Geosciences, Rechbauerstrasse 12, 8010 Graz, Austria (dorothee.hippler@tugraz.at)

²University of Bremen, Institute of Geosciences, Klagenfurter Straße, 28359 Bremen, Germany (a.kluegel@uni-bremen.de)

³Center for Environmental Remediation, Chinese Academy of Sciences, Beijing 100101, China (guojq@igsnr.ac.cn)

⁴Technical University Berlin, Department of Mineralogy, Ackerstrasse 76, 13355 Berlin, Germany (gerhard.franz@tu-berlin.de)

The Precambrian-Cambrian time interval represents one of the greatest phosphogenic episodes in Earth's history with giant and well-preserved phosphate deposits occurring on the Yangtze Platform (South China). We investigated concentrations of rare earth elements (REE) and yttrium of shallow and deep-water sedimentary phosphate deposits of the Ediacaran Doushantuo Formation and the Early Cambrian Zhongyicun Formation by using LA-ICP mass spectrometry. The aim is to examine the temporal and spatial variability of seawater chemistry in conjunction with the conditions of phosphate formation and the evaluation of the extent of diagenetic modification. The REE+Y patterns of shallow-water phosphate deposits exhibit the evolution from flat shale-like to gently inclined seawater-derived patterns, with the Early Cambrian phosphate deposits revealing distinct negative Ce- and positive Y-anomalies indicative for oxygenated surface waters. REE+Y patterns of phosphate deposits of the deep-water facies are flat to highly enriched in MREE, which is manifested in variably pronounced concave-down patterns. In detail, the patterns display different Ce-anomalies, as well as small positive Eu-anomalies. We propose that REE+Y patterns of Ediacaran and Early Cambrian sedimentary phosphate deposits can inherit both primary and secondary signatures reflecting either seawater composition or diagenetic modification and fluid flow. The combination of imaging techniques and in-situ LA-ICPMS thereby enables a more sophisticated examination of the potential sources and processes than whole rock determinations. Placing the results in stratigraphic order and assuming the Ce-anomaly to be a proxy for seawater composition and oxygenation, analog to what is known for carbonates, our results point to evolving depositional niches with surface waters enriched in oxygen.