

Water distribution in the continental and oceanic upper mantle

ANNE H. PESLIER¹, LUC S. DOUCET², MICHAEL BIZIMIS³, HEJIU HUI⁴ AND LILLIAN S. SCHAFFER⁵

¹Jacobs, NASA Johnson Space Center, Mailcode XI3, Houston, TX 77058, USA; anne.h.peslier@nasa.gov;

²Laboratoire G-Time, Université Libre de Bruxelles, Belgium;

³Earth and Ocean Sciences, University of South Carolina, USA;

⁴Earth Sciences and Engineering, Nanjing University, China;

⁵Earth and Atmospheric Sciences, University of Houston, USA.

Nominally anhydrous minerals such as olivine, pyroxene and garnet can accommodate tens to hundreds of ppm H₂O in the form of hydrogen bonded to structural oxygen in lattice defects [1]. Although in seemingly small amounts, this water can significantly alter chemical and physical properties of the minerals and rocks (e.g., [2-4]). Water in particular can modify their rheological properties and its distribution in the mantle derives from melting and metasomatic processes and lithology repartition (pyroxenite vs peridotite) [5-7]. These effects will be examined here using Fourier transform infrared spectrometry (FTIR) water analyses on minerals from mantle xenoliths from cratons, plume-influenced cratons and oceanic settings.

In particular, our results on xenoliths from three different cratons [5] [6] [8] will be compared. Each cratonic mantle lithosphere (Kaapvaal, Siberian and Slave) has a different water concentration distribution that is clearly linked to local metasomatic processes. Only the mantle root beneath the Kaapvaal (southern Africa) has evidence for pervasive dry olivine (< 5 ppm H₂O) at its base (> 6 GPa). This challenges the link between olivine water content and survival of Archean cratonic mantle, and questions whether xenoliths are representative of the whole cratonic mantle. We will also present our latest data on Hawaii and Tanzanian craton xenoliths [7] [9] [10] which both suggest the intriguing result that mantle lithosphere is not enriched in water when it interacts with melts from deep mantle upwellings (plumes).

[1] Bell & Rossman (1992) *Science* **255**, 1391-1397. [2] Chopra & Paterson (1984) *JGR* **89**, 7861-7876. [3] Green (1973) *EPSL* **19**, 37-45. [4] Karato (1990) *Nature* **347**, 272-273. [5] Peslier *et al.* (2010) *Nature* **467**, 78-81. [6] Peslier *et al.* (2012) *GCA* **97**, 213-246. [7] Bizimis & Peslier (2015) *CG* **397**, 61-75. [8] Doucet *et al.* (2014) *GCA* **137**, 159-187. [9] Peslier & Bizimis (2015) *GGG* **accepted**. [10] Hui *et al.* (2015) *GGG* **in revision**.