

## Hf and Nd isotope systematics of early Archean komatiites from the Barberton Greenstone Belt, South Africa

JANNE BLICHERT-TOFT<sup>1</sup>, ALLAN WILSON<sup>2</sup>, AND NICHOLAS T. ARNDT<sup>3</sup>

<sup>1</sup>ENS Lyon, 69007 Lyon, France (jblicher@ens-lyon.fr)

<sup>2</sup>University of the Witwatersrand, Johannesburg, South Africa

<sup>3</sup>Université Grenoble Alpes, 38041 Grenoble, France

To constrain the origin of komatiites from the Barberton Greenstone Belt, South Africa, we measured <sup>176</sup>Lu-<sup>176</sup>Hf and <sup>147</sup>Sm-<sup>143</sup>Nd compositions for 18 komatiites from ICDP coring in the Komati Formation of the Barberton Belt, as well as 33 komatiites from surface outcrops of the Komati, Hooggenoeg, and Weltevreden Formations. Magmatic clinopyroxene from surface samples near the drill site was also analyzed. For the Lu-Hf isotope system, the Komati Formation surface and core samples and the clinopyroxene define a linear array whose slope corresponds to an age of  $3426 \pm 16$  Ma (MSWD = 118), which is slightly younger than the accepted age of the rocks (3.48 Ga). The Sm-Nd isotope data for the same set of samples likewise fall along a linear array also yielding a younger age of  $3339 \pm 12$  Ma (MSWD = 42). The high MSWD for both isotope systems indicate substantial scatter at variance with normal magmatic processes, likely implying element mobility disturbing even these relatively robust isotopic systems shortly after eruption of the lavas. The average initial  $\epsilon_{\text{Hf}}$  and  $\epsilon_{\text{Nd}}$  of the core samples are +1.4 and +0.45, respectively, in accordance with a depleted mantle source at 3.5 Ga, while the clinopyroxene and their host rocks have strongly positive  $\epsilon_{\text{Hf(T)}}$  of about +5 and negative  $\epsilon_{\text{Nd(T)}}$  of about -2. Given the overall scatter of the whole-rock data, the most robust constraint on the composition of the komatiite source comes from the clinopyroxene. Their positive  $\epsilon_{\text{Hf(T)}}$  is in line with, though somewhat higher than other results from komatiites from the Komati Formation, but their negative  $\epsilon_{\text{Nd(T)}}$  is unexpected in that it indicates a source with long-term low Sm/Nd, which is at odds with its long-term high Lu/Hf. This signature is also found in the trace element compositions of some of the komatiites, such as moderately enriched LREE, negative Hf anomalies, and low Hf/Sm ratios. The origin of these features is uncertain but one possibility is that the discordance between the Hf and Nd isotope systems reflects the presence of deep-sea sediments in the source of some of the Barberton komatiites. This has potentially wide implications for plate tectonics and crust-mantle interaction in the early Earth