

Investigation of Cesium adsorption on calcareous soil combined Batch and EDS approaches

JIANJUN LIANG,¹ ZHENGQUAN YAN,² LEI HU,²
XIAOLAN ZHAO,¹ QIAOHUI FAN,¹

¹Key Laboratory of Petroleum Resources Research,
Institute of Geology and Geophysics, CAS,
Lanzhou, 730000 CHINA

²School of Material and Chemical Engineering, West
Anhui University, Lu'an 237012, CHINA

Since last century, a large amount of radiocesium (RCs) released from atomic weapon tests and nuclear accidents was directly introduced into environment through atmospheric transportation and deposition on land surface soil, discharged into river system by erosion effect during rainfall, and finally released into ocean. In this study, the generalized adsorption model (GAM) for Cs⁺ was employed to estimate the solid-water distribution of Cs⁺ in the river-estuary-ocean system. The results confirmed that the capacity of each adsorption sites of river sediments, i.e., frayed edge site, interlayer site, type II site, or planar site, can be precisely optimized through the adsorption isotherm of Cs⁺ combined with the radiocesium interception potential (*RIP*) and cation exchange capacity (*CEC*).

According to the GAM, the main contributor to Cs⁺ adsorption is the frayed edge site due to the very low Cs⁺ concentration in the river-estuary-ocean system. The different solid-water distribution of Cs⁺ in the river-estuary-ocean system was dominantly controlled by the salinity in the aqueous phase. Therefore, cesium should be highly reactive with strong adsorptive character to particulate matters in river system, whereas a conservative distribution must be dominant in ocean with much weaker affinity to particulate matters because of the high salinity.