

## Using meteorological ensembles for atmospheric dispersion modelling of the Fukushima nuclear accident

R. PERILLAT<sup>1,2</sup>, I. KORSAKISSOK<sup>1</sup>, V. MALLET<sup>3</sup>,  
A. MATHIEU<sup>1</sup>, T. SEKIYAMA<sup>4</sup>, M. KAJINO<sup>4</sup>, K.  
ADACHI<sup>4</sup>, T. MAKI<sup>4</sup>, Y. IGARASHI<sup>4</sup>, D. DIDIER<sup>1</sup>

<sup>1</sup>IRSN, PRP-CRI, SESUC, BMCA, avenue de la  
Division Leclerc, B.P. 17, 92262 Fontenay-aux-  
roses, France

(correspondance\* : irene.korsakissok@irsn.fr,  
raphael.perillat-bertintechnologies@irsn.fr,  
anne.mathieu@irsn.fr, damien.didier@irsn.fr)

<sup>2</sup>BERTIN Technologie, 25 route de l'Orme,  
91190 Saint-Aubin, France

<sup>3</sup>INRIA, Clime, 2 rue Simone Iff, 75012 Paris, France  
(vivien.mallet@inria.fr)

<sup>4</sup>MRI, Meteorological Research Institute, Tsukuba,  
Japan (tsekiyam@mri-jma.go.jp, kajino@mri-  
jma.go.jp, adachik@mri-jma.go.jp, tmaki@mri-  
jma.go.jp, yigarash@mri-jma.go.jp)

Dispersion models are used in response to an accidental release of radionuclides of the atmosphere, to infer mitigation actions, and complement field measurements for the assessment of short and long term environmental and sanitary impacts. However, the predictions of these models are subject to important uncertainties, especially due to input data, such as meteorological fields or source term (Korsakissok et al., 2013, Girard et al., 2014).

In the framework of the SAKURA project, an MRI-IRSN collaboration, a meteorological ensemble of 20 members designed by MRI (Sekiyama et al. 2015), designed to reduce the meteorological uncertainty in the Fukushima case, was used with IRSN's atmospheric dispersion models.

First, it was necessary to assess the quality of the ensembles for our purpose, to ensure that their spread was representative of the uncertainty of meteorological fields by using meteorological observations. Then, the uncertainty was propagated through atmospheric dispersion models. The underlying question is whether small uncertainties in meteorological fields can produce large differences in atmospheric dispersion results. In the case of the Fukushima accident, gamma dose rates, air activities and deposition data were available. Based on these data, selection criteria for the ensemble members were designed. Finally, the total uncertainty, including source term and model parameterization, was propagated through the model. Results were compared with the meteorological-induced uncertainty, and between another set of meteorological data retrieved from ECMWF.