

## Evidence for Archean hydrous deep-mantle reservoir provided by Abitibi komatiites

A.V. SOBOLEV<sup>1,2</sup>, E. V. ASAFOV<sup>2</sup>, A. A. GURENKO<sup>3</sup>, N. T. ARNDT<sup>1</sup>, V.G. BATANOVA<sup>1,2</sup>, M.V. PORTNYAGIN<sup>2,4</sup>, DIETER GARBE-SCHÖNBERG<sup>5</sup> AND S. P. KRASHENINNIKOV<sup>2</sup>

<sup>1</sup> Université Grenoble Alpes, ISTERre, Grenoble, France; alexander.sobolev@univ-grenoble-alpes.fr

<sup>2</sup> Vernadsky Institute, Moscow, Russia; evasafov@gmail.com

<sup>3</sup> CRPG, Nancy, France, agurenko@crpg.cnrs-nancy.fr

<sup>4</sup> GEOMAR, Kiel, Germany; mportnyagin@geomar.de

<sup>5</sup> CAU Kiel Uni, Germany; d.garbeschoenberg@gmx.de

Archean komatiites result from melting under extreme conditions of the Earth's mantle. Their chemical compositions evoke very high eruption temperatures, up to 1600°C, providing clues to still higher temperatures in their mantle source [1]. This message is clouded, however, by uncertainty about the water content in komatiite magmas. One school holds that komatiites were essentially dry and originated in mantle plumes [2] while the other argues that these magmas contained several percent of water, which drastically reduced their eruption temperature and links them to subduction processes [3].

We report contents of water and other volatile components, major and trace elements in melt inclusions in olivine (up to Fo 94.5 mol%) and direct estimates of the composition and crystallization temperature of the parental melts of 2.7 Ga old komatiites from the Abitibi belt, Canada [4]. We show that their parental melt contained 30 wt.% MgO 0.6 wt.% H<sub>2</sub>O, and yielded a potential temperature of 1725± 50°C. This melt began to crystallize at around 1510°C at shallow depth and under reducing conditions (QFM-1), and it evolved via fractional crystallization of olivine, accompanied by minor crustal assimilation. As its major and trace element composition are inconsistent with a subduction setting, we propose that the unusually high ratio of H<sub>2</sub>O to elements of similar compatibility (H<sub>2</sub>O/Ce > 6000) was caused by entrainment into the komatiite source of hydrous material from the mantle transition zone [5]. These results confirm a plume origin of komatiites and high Archean mantle temperatures, and evoke a hydrous reservoir in the deep mantle early in the Earth history.

[1] Herzberg, C. *JGR* **97**, 4521-4540 (1992). [2]. Arndt, N. et al. *Geology* **26**, 739-742, (1998). [3] Grove, T. L. & Parman, S. W. *EPSL* **219**, 173-187, (2004). [4]. Sobolev et al, *Nature* **531**, DOI 10.1038/nature17152 (2016). [5]. Bercovici, D. & Karato, S. *Nature* **425**, 39-44 (2003).