

Making Mercury's Core with Light Elements

KATHLEEN E. VANDER KAADEN*^{1,2}, FRANCIS M. MCCUBBIN^{1,2}, D. KENT ROSS^{2,3,4}

¹Institute of Meteoritics, Department of Earth and Planetary Sciences, 1 University of New Mexico, MSC03-2050, Albuquerque, NM 87131.

²NASA Johnson Space Center, Mail Code XI2, 2101 NASA Parkway, Houston, TX 77058

³Jacobs JETS, NASA Johnson Space Center, 2101 NASA Parkway, Houston TX 77058

⁴University of Texas at El Paso-CASSMAR

(*correspondence:kathleen.e.vanderkaaden@nasa.gov)

Recent results obtained from the Mercury Surface, Space ENvironment, GEochemistry, and Ranging spacecraft showed the surface of Mercury has low FeO abundances (<2 wt%) and high S abundances (~4 wt%) [1], suggesting the fO_2 of Mercury's surface materials is somewhere between 3 to 7 \log_{10} units below the IW buffer [2]. The highly reducing nature of Mercury has resulted in a relatively thin mantle and a large core that has the potential to exhibit an exotic composition in comparison to the other terrestrial planets. This exotic composition may extend to include light elements (e.g., Si, C, S). Furthermore, [3] has argued for a possible primary floatation crust on Mercury composed of graphite, which may require a core that is C-saturated. In order to investigate mercurian core compositions, we conducted piston cylinder experiments at 1 GPa, from 1300 °C to 1700 °C, using a range of starting compositions consisting of various Si-Fe metal mixtures (Si_5Fe_{95} , $Si_{10}Fe_{90}$, $Si_{22}Fe_{78}$, and $Si_{35}Fe_{65}$). All metals were loaded into graphite capsules used to ensure C-saturation during the duration of each experimental run. Our experiments show that Fe-Si metallic alloys exclude carbon relative to more Fe-rich metal. This exclusion of carbon commences within the range of 5 to 10 wt% Si. These results indicate that if Mercury has a Si-rich core (having more than ~5 wt% silicon), it would have saturated in carbon at low C abundances allowing for the possible formation of a graphite floatation crust as suggested by [3]. These results have important implications for the thermal and magmatic evolution of Mercury. **References:** [1] Nittler, L.R. *et al.*, (2011) *Science* [2] McCubbin, F.M. *et al.*, (2012) *GRL*. [3] Vander Kaaden, K.E. and McCubbin, F.M. (2015) *JGR-Planets*.