

Copper Isotope Metallomics and Amyotrophic Lateral Sclerosis

T. GABRIEL ENGE^{1*}, ANTHONY DOSSETO¹,
HEATH ECROYD², DIANNE F. JOLLEY³

¹Wollongong Isotope Geochronology Laboratory,
School of Earth and Environmental Sciences,
University of Wollongong, NSW, Australia
(*correspondence: tge571@uowmail.edu.au;
tonyd@uow.edu.au)

²Illawarra Health and Medical Research Institute,
University of Wollongong, NSW, Australia
(heathe@uow.edu.au)

³School of Chemistry, University of Wollongong,
NSW, Australia (djolley@uow.edu.au)

Amyotrophic lateral sclerosis (ALS) is a devastating neurodegenerative disease in which motor neurons undergo progressive degeneration and die. Evidence indicates that ALS is a protein misfolding disorder [1]. Some forms of familial ALS are due to mutations in the Cu, Zn-superoxide dismutase 1 (SOD1) protein which destabilise it, causing it to misfold and accumulate into deposits within the cells. A reason why these mutations destabilise SOD1 could be that the binding of Cu by SOD1 may become defective [2].

An early diagnostic biomarker for ALS is essential in developing future potential treatment options, since the pathology develops well before symptoms do. Our work uses a transgenic mouse model of ALS to evaluate whether changes to Cu metabolic pathways and metal concentrations in specific tissues are associated with the onset and progression of ALS and could be used as a biomarker of disease. We have collected tissue samples from transgenic mice (n=18) at key points during disease progression (30, 60, 90 and 120 days) as well as healthy controls (n=18) and measured the Cu isotope composition ($\delta^{65}\text{Cu}$) and Cu, Zn, Ni, Co and Fe concentrations in brain, spinal cord, whole blood and muscle tissue (quadriceps femoris).

Our results demonstrate that Cu, Fe and Zn concentrations in the muscle tissue of ALS mice are significantly higher than in healthy mice. This is also the case for Cu in spinal cord. While there is no difference in Cu concentration in blood between these two groups, preliminary results for Cu isotopes suggest differences in the isotopic composition of blood between ALS and healthy mice, warranting assessment if blood could be used as an early disease-biomarker. We will present additional results which should shed light on the links between Cu isotope metallomics and ALS disease progression.

- 1] Broom, *et al.* (2014) *Essays Biochem* **56**, 149-65.
[2] Hayward, *et al.* (2002) *J. Biol Chem* **227**, 15923-31.