

Deep cycle of sulphur: the role of subducted sediments

A.-A. PELLETER^{1*}, G. PROUTEAU¹, B. SCAILLET¹

¹Univ. Orléans, ISTO/CNRS, UMR 7327, 45071 Orléans, France; BRGM, UMR 7327, BP 36009, 45060 Orléans, France.

(*correspondence: anne-aziliz.pelleter@univ-orleans.fr)

It is widely accepted that sediment-derived silicate melts are required to efficiently transfer key trace elements (Be, Th) towards the mantle wedge [1,2]. If recent experimental studies highlight the role of H₂O, CO₂ or halogens (Cl, F) during the partial melting of subducted sediments, effect of sulphur remains poorly constrained [3]. Yet, this volatile element is often mentioned as a potential oxidising agent of the mantle wedge (as S⁶⁺) [4] and geochemical studies show that sub-arc mantle is significantly enriched in S relative to MORB mantle following the percolation of a metasomatic agent expelled from subducted lithologies [5]. The main purpose of this study is to evaluate the ability of different sediment-derived silicate melts to transfer sulphur in the sub-arc mantle.

Using piston-cylinder apparatus, we performed melting experiments (3 GPa ; 750-1000°C ; $f_{O_2} \sim NNO-NNO+1$) on two types of natural, trace-element undoped and volatile-rich sediments: (1) a Ca-poor pelite and (2) a marlstone. These compositions correspond to the two main endmembers of sediments subducted worldwide. For the purpose of this study, some experiments were doped with elemental sulphur (1 and 2 wt%).

Silicate melts produced by the fluid-present melting of the pelite range from trondhjemitic to granitic compositions and are broadly peraluminous (A/CNK until 3.02). S contents in these melts range between 72 to 1033 ppm with a maximum at 850-900°C. For fluid-present melting of the marlstone, produced silicate melts are granodioritic and metaluminous or slightly peraluminous ($0.95 < A/CNK < 1.07$). S contents range from 371 to 2985 ppm and increase with the temperature. Preliminary results (raman spectroscopy) show that S can be dissolved as SO₄²⁻ (S⁶⁺) in silicate melts derived from the pelitic sediment.

[1] Johnson & Plank (1999) *Geochem. Geophys. Geosyst.* **1**, 1007. [2] Hermann & Rubatto (2009) *Chem. Geol.* **265**, 512-526. [3] Prouteau & Scaillet (2013) *J. Petrol.* **54**, 183-213. [4] Kelley & Cottrell (2009) *Science* **325**, 605-607 [5] Wallace & Edmonds (2011) *Reviews in Mineralogy and Geochemistry* **73**, 215-246.