

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

## $\delta^{26}\text{Mg}$ record of Phanerozoic oceans

P. B. TÖRBER<sup>1,2,3\*</sup>, J. FARKAŠ<sup>2,1,3</sup>, C. ROLLION-BARD<sup>4</sup>, K. WALLMANN<sup>5</sup>, U. BRAND<sup>6</sup>, K. AZMY<sup>7</sup>, A. TOMAŠOVÝCH<sup>8</sup>, C. LÉCUYER<sup>9</sup>, N. VIGIER<sup>10</sup>, S. SAULNIER<sup>11</sup>, M. KOMÁREK<sup>1</sup>, T. MAGNA<sup>3</sup>, M. ŠIMEČEK<sup>3</sup>, M. FRANCOVÁ<sup>3</sup>, F. BÖHM<sup>5</sup>, AND A. EISENHAEUER<sup>5</sup>

\*Email (corresponding author): toerber@fzp.czu.cz

<sup>1</sup>Czech University of Life Sciences Prague, Czech Republic

<sup>2</sup>University of Adelaide, Australia; <sup>3</sup>Czech Geological Survey, Czech Republic; <sup>4</sup>CNRS/IPGP Paris, France;

<sup>5</sup>GEOMAR, Germany; <sup>6</sup>Brock University, Canada; <sup>7</sup>Memorial University of Newfoundland, Canada; <sup>8</sup>Slovak Academy of Sciences, Slovakia; <sup>9</sup>University of Lyon, France;

<sup>10</sup>Villefranche Oceanographic Laboratory, France;

<sup>11</sup>Laboratory of Physical Geography, France

The secular variation in the marine Mg/Ca ratio over geologic time is undisputed, however, the role of driving forces behind this phenomenon remains uncertain. A key to the discrimination of major fluxes is the quantification of the Mg oceanic cycle that in turn can be examined by using the Mg isotope compositions of low-Mg calcite brachiopod shells. Within the BASE-LiNE Earth project, a total of 95 analyses on modern and fossil (early Ordovician to Quaternary) brachiopod specimens were performed to generate an inferred  $\delta^{26}\text{Mg}$  paleo-seawater record. As a prerequisite, selected modern, globally distributed species, whose average “habitat temperatures” range from about 0 to 29°C, were investigated and results revealed a weak ( $\sim 0.02\text{‰}\text{°C}^{-1}$ ) temperature-sensitivity of  $\delta^{26}\text{Mg}$  in shells. This supports their suitability for paleo-seawater  $\delta^{26}\text{Mg}$  reconstructions. The offset  $\Delta^{26}\text{Mg}$  between modern global seawater  $\delta^{26}\text{Mg}$  [1] and the average of modern brachiopods is about  $-1.26\text{‰}$ , and has been applied to fossil samples. A preliminary locally weighted and smoothed  $\delta^{26}\text{Mg}$  paleo-seawater trend yielded (i) short-term negative and positive anomalies during the Cenozoic, (ii) rather constant values during most of the Mesozoic, (iii) a significant positive-to-negative shift during the Permian/Carboniferous transition, and (iv) systematically negative values during the rest of the Palaeozoic (relative to modern seawater). This composite  $\delta^{26}\text{Mg}$  record of Phanerozoic seawater will be simulated via a coupled numerical model of oceanic elemental cycles [2], and conclusions will be made regarding the plausible driving mechanism(s) behind the observed long-term changes in the marine Mg/Ca record.

[1] Ling et al. (2011) *Rapid Commun. Mass Spectrom.* **25** 2828–2836. [2] K. Wallmann (2001), *Geochim. Cosmochim. Acta* **65**, 3005–3025.