

Compound-specific degradation of phosphorus sources
by the marine diatoms *Thalassiosira* spp.

ALISIA HOLLAND¹, JAMES G. SANDERS¹, DOUGLAS
MOLLETT^{1,2}, SOLANGE DUHAMEL³, YUANZHI TANG⁴,
JULIA M. DIAZ¹

¹ Skidaway Institute of Oceanography, Department of Marine
Sciences, University of Georgia, Savannah, Georgia, USA

² Georgetown College, Georgetown Kentucky, USA
(drmollett@georgetowncollege.edu)

³ Lamont-Doherty Earth Observatory of Columbia University,
Palisades, New York, USA
(sduhamel@ldeo.columbia.edu)

⁴ School of Earth and Atmospheric Sciences, Georgia
Institute of Technology, Atlanta GA, USA
(yuanzhi.tang@eas.gatech.edu)

Phosphorus (P) is an essential nutrient that can limit ocean productivity and the biological carbon dioxide (CO₂) pump over a range of timescales. In the marine environment, microbial communities recycle bioavailable P by employing enzymes known as phosphatases to degrade complex P-containing molecules under a range of prevailing nutrient concentrations, from oligotrophic to eutrophic. Here, we examined the transformation of various P-containing molecules by representative diatoms of the genus *Thalassiosira*. Rates of orthophosphate production arising from the degradation of complex P species, as well as proteomic analysis of phosphatase enzyme diversity, revealed preferential degradation of P species containing phosphoanhydride (P-O-P) bonds, such as inorganic polyphosphate. The transformations of these diverse P-containing species by diatoms has implications beyond nutrient cycling, as the degradation of polyphosphate has previously been linked to the sequestration of P via calcium phosphate mineral precipitation. Therefore, this work may act to clarify the role of marine microorganisms in shaping long-term inventory of marine P, and consequently illuminate feedbacks on global marine primary productivity, ecosystem health, and climate.