

Mineral colonization samplers reveal patterns in microbial community composition and structure at hydrothermal vents

HEATHER OLINS¹ AMY GARTMAN² PETER GIRGUIS³

¹Department of Biology, Boston College, Chestnut Hill, MA
olins@bc.edu

²US Geological Survey, Santa Cruz, CA

³Department of Organismic & Evolutionary Biology,
Harvard University, Cambridge, MA

Microbes can use solid phase minerals as both electron donors and acceptors, and thus minerals may exert a selective pressure on microbes (especially some less common taxa) colonizing their surfaces. Little is known about the extent of this selection. Hydrothermal vent chimneys are ideal to investigate these microbe-mineral interactions. In order to examine the extent to which solid-state minerals influence community structure in hydrothermal vent chimneys, we designed and deployed novel *in situ* mineral associated colonization samplers (ISMACS), with which we incubated a variety of minerals in vent fluid. Gross community composition did not vary significantly with mineralogy. Certain taxa did appear to be significantly more abundant on specific minerals. We found evidence that communities exposed to a mix of mineral substrates were more diverse than single-mineral communities, and that hematite-hosted communities were less diverse than other minerals. These data indicate that mineral substrate is not the primary driver of community structure in hydrothermal vent chimneys, but that it plays an important role in the diversity, succession, and metabolism of these microbial communities – especially in rare taxa.