

Seamount subduction at seismogenic depths: Structural and metamorphic evidence from the Zagros suture zone

BONNET G.¹, AGARD P.², ANGIBOUST S.³, FOURNIER M.
⁴ OMRANI J.⁵

¹ Sorbonne Université, 4 place Jussieu, 75005 Paris, France,
(gbonnet01@gmail.com)

² Sorbonne Université, 4 place Jussieu, 75005 Paris, France,
(philippe.agard@upmc.fr)

³ Sorbonne Paris Cité, 1 place Jussieu, 75005 Paris, France,
(angiboust@ipgp.fr)

⁴ Sorbonne Université, 4 place Jussieu, 75005 Paris, France,
(marc.fournier@upmc.fr)

⁵ Geological Survey of Iran, Mehraj Boulevard, Tehran, Iran
(j.omrani@gmail.com)

Large-scale seafloor topographic features, such as seamounts, are for the most part subducted with the downgoing oceanic plate. They are expected to critically impact the seismogenic and mechanic behavior of subduction zones, but their exact role is strongly debated (i.e., as to whether they represent barriers to propagation or asperities promoting nucleation).

Rare natural examples of metamorphosed seamounts, that got sliced off the slab along the plate interface and escaped recycling into the mantle, are therefore precious witnesses to document processes operating at depths of 0-30 km.

We herein report the existence of a large-scale oceanic topographic structure sandwiched in the Zagros suture zone (Siah Kuh unit), most probably a former seamount, along with other blueschist units (Angiboust et al., EPSL 2016). The main criteria for identifying this seamount are its shape, lithologies, sedimentary cover and geochemistry.

HP-LT minerals (lawsonite, aragonite veins, blue amphibole) found across the whole structure, particularly in zones of localized compressive deformation, indicate that this seamount was shallowly subducted at ~20 km. This deformation, interpreted to be syn-subduction, is assisted by décollement rooting in serpentinite and/or oceanic metasediments and is associated with rare cataclase in magmatic rocks. We interpret these structures as related to the internal slicing of the seamount in subduction. The presence of these soft layers may prevent seismogenic deformation, since no pseudotachylites have been found. The Siah Kuh unit is also a perfect target to investigate subduction processes such as mechanical coupling, fluid/tectonic overpressure and the nature of subduction fluids.

**This abstract is too long to be accepted for publication.
Please revise it so that it fits into the column on one
page.**