

**Am(III)/Nd(III) INTERACTIONS
WITH BORATE:
EXPERIMENTAL
INVESTIGATIONS OF
Nd(OH)₃(micro cr) SOLUBILITY IN
MIXTURES of NaCl AND MgCl₂ IN
EQUILIBRIUM WITH BORAX ^A**

YONGLIANG XIONG^{1*}, LESLIE KIRKES¹, JANDI KNOX,
CASSIE MARRS¹, HEATHER BURTON¹

¹*Sandia National Laboratories (SNL)^B, Carlsbad Programs
Group, 4100 National Parks Highway, Carlsbad, NM
88220, USA*

**Corresponding author email: yxiong@sandia.gov*

In previous studies on Am(III)/Nd(III) interactions with borate, the borate concentrations were kept at a relatively low level, up to $\sim 0.16 \text{ mol}\cdot\text{kg}^{-1}$ as the highest. Such concentrations of borate are not saturated, in terms of borate, with borate-bearing phases such as borax ($\text{Na}_2\text{B}_4\text{O}_7\cdot 10\text{H}_2\text{O}$) and tincalconite ($\text{Na}_2\text{B}_4\text{O}_7\cdot 5\text{H}_2\text{O}$) that could be present in geological repositories. Therefore, the impact of Am(III)/Nd(III) interactions with borate is not fully understood. For instance, tincalconite has been observed as a corrosion product for borosilicate glass for HLW under the repository conditions in China.

In this study, we investigate the Nd(III) interactions with borate in equilibrium with borax in the mixtures of NaCl and MgCl₂ at 25°C, saturated with borax. In our experiments, the solubility-controlling phase for Nd(III) is Nd(OH)₃(micro cr). In synthesis of Nd(OH)₃(micro cr), we followed the procedure from literature. High purity Nd₂O₃ was loaded into Paar® reaction vessels with deaerated DI water. Nd(OH)₃(micro cr) formed when the high purity Nd₂O₃ reacted with the deaerated DI water at 200°C in Paar® reaction vessels for a period of two weeks.

Our results indicate that high borate concentrations ($\sim 0.5 \text{ mol}\cdot\text{kg}^{-1}$) have an obvious effect on Nd(III), enhancing the solubility of Nd(OH)₃(micro cr).

^A This research is funded by the WIPP programs administered by the Office of Environmental Management (EM) of the U.S. Department of Energy.

^B Sandia National Laboratories is a multimission laboratory operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA-0003525. SAND2018-3288A