

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

## **Laboratory experimental study of CO<sub>2</sub>-brine-synthetic sandstone core interaction under high P/T reservoir conditions**

ZHICHAO YU<sup>1,2\*</sup>, KEYU LIU<sup>3</sup>, LILIU<sup>4</sup>

<sup>1</sup> Research Institute of Petroleum Exploration & Development, PetroChina, Beijing, 100083, P.R. China  
Yuzhichao@petrochina.com.cn(\*)

<sup>2</sup> Key Laboratory of Basin Structure & Hydrocarbon Accumulation, CNPC, Beijing, 10083, P.R. China

<sup>3</sup> School of Geosciences, China University of Petroleum, Qingdao, Shandong, P.R. China  
keyu.liu@icloud.com

<sup>4</sup> College of Earth Sciences, Jilin University, Changchun 130061, P.R. China  
liuli0892@vip.sina.com

To understand the interaction of CO<sub>2</sub> and sandstone after CO<sub>2</sub> injection under ultra-pressure reservoir conditions, we carried out CO<sub>2</sub>-brine-rock experiments using a batch stainless steel reactor under high pressure/temperature reservoir conditions (150°C and 48 MPa, pressure coefficient=1.5). Six synthetic sandstone cores having the same mineral constituents with the real reservoir sandstone are used in the experiment.

Changes in the ionic chemistry of the outlet solution, combined with core scanning electron microscopy (SEM) analysis of the core pre- and post-experiments reveal new insights into CO<sub>2</sub>-brine-rock interactions. Minerals such as quartz, potassium (K) feldspar, albite, dolomite and calcite are variably dissolved after the experiments. Calcite is the mineral mostly affected by dissolution, secondly by dolomite, followed by the feldspar minerals, while dissolution of quartz is minimal. Secondary porosity increased gradually with reaction time, whereas the primary porosity decreased. Core porosity increased substantially after the experiment. The increase in the porosity is primarily from the dissolution of carbonate minerals. No CO<sub>2</sub> sequestration minerals were precipitated after the experiment. Precipitation of carbonates related to CO<sub>2</sub> injection may have been inhibited if the solution did not attain the critical saturation state required for nucleation. The results provide some new insights into CO<sub>2</sub> trapping mechanisms in high pressure/temperature reservoirs, and into the understanding of geochemical reactions in CO<sub>2</sub>-brine-rock systems that is particularly relevant to the geologic CO<sub>2</sub> sequestration.

*Keywords: CO<sub>2</sub> sequestration; dissolution; precipitation*