

***In situ* LA-ICPMS U-Pb dating of fluorite**

THOMAS BLAISE¹, LOUISE LENOIR¹, FREDERIC HAURINE¹, BENJAMIN BRIGAUD¹, JOCELYN BARBARAND¹, MAURICE PAGEL¹

¹ Géosciences Paris Sud (GEOPS), Univ. Paris-Sud, CNRS, 91405 Orsay Cedex, France
(correspondence: thomas.blaise1@u-psud.fr)

Unconformity-related ore deposits are often resulting from multiple fluid flow events generating mineral dissolution-recrystallization, remobilization and precipitation [1,2]. Dating crystallization events in such complex systems is a challenge, especially in the absence of significant U-bearing phases. Fluorite Sm-Nd geochronology may provide robust age estimates, but errorchrones and imprecise ages may occur. The recent development of *in situ* U-Pb radiometric dating of common Pb-bearing minerals such as calcite and hematite using LA-ICPMS offered the exciting possibility to precisely date multiple crystallization events in ore deposits [1].

Here, we present preliminary results on fluorite U-Pb dating from the world-class Pierre-Perthuis unconformity-related F-Ba ore deposit (southeast of the Paris Basin, France). Such research was motivated by the presence of a zonation in fluorite that cannot be individually sampled for Sm-Nd geochronology, and by the absence of calcite or other datable authigenic minerals. Fluorite crystals display a ca. 200µm-thick external growth band in which U concentration varies between 1 to 10 ppm. We have sampled and analyzed this U-bearing growth band by LA-ICPMS (Sector Field) to evaluate the applicability of the U-Pb small scale isochron dating method [1] in fluorite [3].

Variable U/Pb ratios result in an excellent spread of data in a Tera-Wasserburg $^{207}\text{Pb}/^{206}\text{Pb}$ vs. $^{238}\text{U}/^{206}\text{Pb}$ diagram. Unanchored linear regression gives a common initial $^{207}\text{Pb}/^{206}\text{Pb}$ ratio of 0.810, MSWD of 1.4, and an age and uncertainty of 33.4 ± 1.6 Ma. Analytical precision and accuracy are limited by variable ablation rates in fluorite and by the absence of matrix-match external standards. However, this raw age appears consistent with a known fluid circulation phase during the opening of the European Cenozoic Rift System (ECRIS). We will discuss efforts to improve precision and accuracy of fluorite U-Pb dating.

[1] Walter B. et al. (2018) GCA 240, 11-42. [2] Chi G. et al. (2018) Econ Geol 113, 1209-1217. [3] Piccione G. et al. (2015) GSA Annual meeting abstract, vol. 47, No. 7, p. 379.