

Experimental perspectives on hydrothermal vent fluid chemistry at mid-ocean ridges

W. E. SEYFRIED, JR.^{1*}, P. J. SCHEUERMANN¹,
N. J. PESTER², C. TAN¹

¹Department of Earth Sciences, University of Minnesota,
Minneapolis, MN 55455 (*corresponding wes@umn.edu)

²Department of Earth and Planetary Sciences, University of
California, Berkeley, CA. 94702

Recent advances in the design of reactors for hydrothermal experiments have facilitated investigation of processes relevant to geochemical controls on mass transfer processes in deep sea hydrothermal vent fluids. Thus, experimental data have now been obtained for partitioning of transition metals¹ between coexisting vapor and liquid (isothermal decompression) at unusually challenging chemical and physical conditions. These data reveal surprising increases in Fe/Cl and Mn/Cl in vapor, consistent with similar observations of vent fluid chemistry at EPR 9°N in the aftermath of eruptive events. Further, new experimental data for quartz solubility² at elevated temperature (420-500°C) and low to moderate pressure (31- 51 MPa) in single and two-phase fluids (vapor-liquid) extend the pressure range of density-based algorithms, permitting more accurate estimates of temperature of vent fluids using quartz geothermometry. Applying the quartz solubility data with constraints imposed by phase equilibria in the two-phase NaCl-H₂O system to vapor-rich vent fluid issuing from the Piccard site at Mid-Cayman Rise, suggest temperature and pressures of $540 \pm 15^\circ\text{C}$ and 62.5 ± 3.0 MPa. More recently, pH and redox buffered experiments elucidate the partitioning of dissolved H₂ between coexisting NaCl-bearing vapor and liquid, with quantitative implications for establishing subsurface redox controls from dissolved H₂ in hydrothermal vent fluids. Combination of experimental data with time series observations of vent fluid chemistry and data from reaction path and reactive transport models provide a particularly effective framework to assess geochemical controls on the temporal evolution of both basalt and ultramafic hosted hydrothermal systems at mid-ocean ridges in time and space.

^[1] Pester and Seyfried (2014) *Geology*. **42**, 255-258.

^[2] Scheuermann, Tan & Seyfried (2018) *Geochemistry, Geophysics, Geosystems* **19**, 3570-3582.