

Potassium stable isotope homeostasis in vertebrates

T. TACAIL^{1*}, J. LEWIS¹, T. TÜTKEN², C. D. COATH¹, N.
LLOYD³, M. CLAUSS⁴, T. ELLIOTT¹

¹Bristol Isotope Group, School of Earth Sciences, University
of Bristol, UK (*theo.tacail@bristol.ac.uk)

²Applied & Analytical Palaeontology, Institut. of Geosciences,
Johannes Gutenberg Univ., Mainz, Germany

³Thermo Fisher Scientific (Bremen) GmbH, Hanna-Kunath
Str. 11, Bremen, Germany

⁴Clinic for Zoo Animals, Exotic Pets and Wildlife, Vetsuisse
Faculty, University of Zurich, Switzerland

Potassium (K) is an essential nutrient to vertebrates, contributing to central functions of cells (*e.g.*, membrane polarisation), tissues (*e.g.*, muscle contraction) and circulatory system (*e.g.*, blood pressure). These functions are maintained through homeostatic mechanisms, either acting on the maintenance of the total body K stores (external homeostasis) or its asymmetric extra-/intracellular distribution (internal homeostasis) in spite of varying environmental and physiological conditions. Such a regulated processing of K by organisms likely induces mass-dependent fractionation of its stable isotopes (⁴¹K, ³⁹K). To date, however, the systematics of the K isotope cycle in vertebrates and its potential as a tool to explore their biology is virtually unexplored.

Here, we present the first results of a systematic study of K isotope distributions in vertebrates (rats, guinea pigs, quails) reared in controlled feeding experiments. All animals were divided in three groups of three individuals each, fed with the same plant-, meat- and insect-based pellets in order to assess trophic and nutritional effects. The K isotope compositions ($\delta^{41/39}\text{K}$, rel. to SRM3141a) were measured in the feed, tissues and fluids (muscle, liver, kidney, bone, blood, urine) using the Thermo Scientific Proteus collision cell MC-ICP-MS/MS and a bespoke purification chemistry.

Our results show that animal soft tissues are globally enriched in heavy isotopes with respect to dietary K (+0.2 to +1.25 ‰), depending on tissues, species and diet. Diet plays a role in determining isotope compositions of animal tissues. However, the variability of tissue $\delta^{41/39}\text{K}$ between and within species groups appear to be mainly controlled by differences in K cycling. We observe significant patterns indicative of mechanisms shared between the three species and likely related to the homeostatic regulation of K stores, notably muscle and liver involved in internal homeostasis.

This work paves the way to the development of an interpretative framework for the future of K isotopes in the study of key aspects of vertebrates biology, such as energy metabolism or pathological disruptions of K homeostasis.