Effects of Benthic Macrofaunal Bioturbation on the Preservation of Diatom Silica and Formation of Authigenic Products

DYLAN COLE¹, NILS VOLKENBORN², AND JEFFREY W. KRAUSE^{1,3}

¹Dauphin Island Sea Lab, Dauphin Island, AL, 36528, USA
(dcole@disl.edu, jkrause@disl.edu*)

²School of Marine and Atmospheric Sciences, Stony Brook University, Stony Brook, NY, 11794, USA
(nils.volkenborn@stonybrook.edu)

³University of South Alabama, Mobile, AL, 36688, USA

Bioturbating macroinfauna can be considered ecosystem engineers; however, the long-term impact of particle reworking and bioirrigation on elemental cycles is not well constrained. We analyzed sediment samples collected from 400 m² experimental plots in the German Wadden Sea, where lugworms (*Arenicola marina*) have been permanently excluded from intertidal sandflats over the last 17 years by a 1-mm polyethylene mesh buried at 10 cm depth in 2002. The mesh prevented adult lugworms from establishing J-shaped burrows which typically are 20-30 cm deep, offering a unique opportunity to investigate the long-term consequences of bioturbation by adult lugworms on sediment silicon cycling. Samples were analyzed for both biogenic silica and acid-leachable silica —the latter being associated with early diagenetic products. Data from two exclusion and two non-exclusion (i.e. control) cores show enhancement in biogenic silica (3–4-fold increases) and acid-leachable silica (2–3-fold increases) in surface sediments (0–5 cm depth) of exclusion plots. For both plot types, biogenic silica was highest at 10-20 cm and declined to minima at 30-40 cm with an increasing proportion of acid-leachable silica with depth, especially in exclusion plots. These data suggest that *A. marina* can significantly modify local silicon cycling by reducing sedimentary retention of biogenic silica and enhancing mobilization and benthic flux of dissolved silicate to the overlying water.