

C-N-He-Ar Cycling at the Hikurangi Subduction Margin, New Zealand

GABE S. EPSTEIN^{1*}, GRAY E. BEBOUT¹, BRUCE W. CHRISTENSON², HIROCHIKA SUMINO³, IKUKO WADA⁴, CYNTHIA A. WERNER⁵, DAVID R. HILTON⁶

¹Dept. of Earth and Environmental Sciences, Lehigh University, Bethlehem, PA, USA (*correspondence: gse216@lehigh.edu)

²National Isotope Center, GNS Science, Lower Hutt, NZ

³Dept. of Basic Science, University of Tokyo, Tokyo, Japan

⁴Dept. of Earth Sciences, University of Minnesota, Minneapolis, MN, USA

⁵USGS-Contractor, New Plymouth, NZ

⁶Scripps Institution of Oceanography, La Jolla, CA USA

We are evaluating the cycling of C, N, and noble gases at the Hikurangi margin, with ~35 analyses of gases from across the forearc-arc-backarc and further analyses planned for April-May 2020. We present C-N concentrations and isotope compositions of sediments outboard of the trench and wall-rock metasediment in the Taupo Volcanic Zone (TVZ). We compare these data with noble gas and C-N data for gases from fumaroles and thermal springs. Ongoing work includes thermal modeling, thermodynamic calculations of prograde devolatilization, and estimation of TVZ CO₂ output flux.

The incoming sediment section at IODP Site 1520 consists of uppermost terrigenous trench-fill (7 ± 3 wt.% carbonate, 0.39 ± 0.17 wt.% organic C), pelagic sediment (61 ± 21 wt.% carbonate, 0.24 ± 0.15 wt.% organic C), and lowermost volcanoclastics (13 ± 14 wt.% carbonate). Isotope compositions are relatively uniform, with δ¹⁵N = +4.4 ± 0.9‰ (AIR), δ¹³C_{carb} = +0.9 ± 1.1‰ (VPDB), and δ¹³C_{red} = -25.9 ± 1.2‰ (VPDB). Wall-rock metasediments have δ¹⁵N = +2.4 to +6.4‰, δ¹³C_{red} = -25.0 ± 1.9‰. Trench-fill sediments are largely removed by accretion, thus the carbonate-rich section likely contributes more to the gas emissions.

The dominant C-bearing gas phase in the forearc is CH₄ (δ¹³C = -35 to -53‰) and that within the TVZ gases is CO₂ (δ¹³C = -2 to -10‰). Forearc noble gas ratios have crustal to atmospheric values (± minor mantle contribution; ³He/⁴He = 0.2-1.7 R_A and ⁴⁰Ar/³⁶Ar ≥ 296), while He-Ar and C-N isotope values of gases from the TVZ are consistent with mantle and recycled sedimentary contributions (³He/⁴He = 4-7 R_A, δ¹⁵N = +1.3 ± 0.9‰, and [N₂/³⁶Ar]/AIR = 1-10). Overlap in δ¹³C_{red} and δ¹⁵N of incoming sediments and wall rocks complicates differentiation of C sources but, given the accretion of the trench-fill sequence, the apparent sediment-derived C_{org} component (≈ 30%, after [1]) and N in the gases could reflect contamination by Torlesse/Waipapa wall rocks.

[1] Sano & Marty (1995) *Chem. Geol.* **119**, 265-274.