

The physical and chemical evolution of magmatic fluids in near-solidus silicic magma reservoirs: Implications for the formation of pegmatites

JULIANA TROCH¹, CHRIS HUBER² AND OLIVIER BACHMANN³

¹Smithsonian National Museum of Natural History

²Brown University

³ETH Zürich

Presenting Author: trochj@si.edu

During their ascent through the crust, magmas cool and differentiate, and water and other fluid-mobile elements (e.g. Li, B, C, F, S, Cl) are continuously concentrated in the residual melt, until reaching fluid saturation. The ensuing exsolution of a magmatic volatile phase (MVP) dominated by H₂O is thought to generally occur early in the evolution of long-lived “mushy” magma reservoirs, and can be simulated by tracking the chemical and physical evolution of these reservoirs in thermomechanical numerical models. Pegmatites are commonly interpreted as the crystallization products of late-stage volatile-rich liquids sourced from granitic igneous bodies. However, little is known about the timing and mechanism of extraction of pegmatitic liquids from their source.

In this study, we review findings from thermomechanical models on the physical and chemical evolution of melt and MVP in near-solidus magma reservoirs and apply these to textural and chemical observations from pegmatites. As an example, we use a 3-phase compaction model of a section of a mushy reservoir, and couple this to fluid-melt and mineral-melt partition coefficients of volatiles trace elements (Li, Cl, S, F, B). We track various physical parameters of melt, crystal and MVP, such as volume fractions, densities, velocities, as well as the content in the volatile trace elements mentioned above. The results suggest that typical pegmatite-like compositions (i.e. enriched in incompatible elements) require high crystallinities (>70-75 vol% crystals) in the magma reservoir, at which MVP is efficiently trapped in the crystal network. Volatile trace elements can become enriched beyond contents expected from closed-system equilibrium crystallization by transport of MVP from more evolved mush domains.

From a thermomechanical perspective, these observations indicate that, rather than from melt, pegmatites may more likely be generated from pressurized, solute-rich MVP with dissolved components of silicate melt and volatile elements. Hydraulic fracturing provides a mechanism for the extraction and emplacement of such pegmatite-generating liquids in and around the main parental near-solidus mush as pockets, dikes, and small intrusive bodies. This thermomechanical framework for the extraction of MVP from mushes, and associated formation of pegmatites integrates both igneous and hydrothermal realms into the concept of transcrustal magmatic distillation columns.