

# Fossil Carbon Release and Exports from Subsurface Sedimentary Bedrock Weathering in Mountainous Watersheds

JIAMIN WAN<sup>1</sup>, TETSU K TOKUNAGA<sup>2</sup>, WENMING  
DONG<sup>2</sup>, CURTIS BEUTLER<sup>3</sup>, ALEXANDER W NEWMAN<sup>3</sup>,  
WENDY BROWN<sup>3</sup>, MARKUS BILL<sup>2</sup>, JENNIFER L  
DRUHAN<sup>4</sup>, AMANDA N HENDERSON<sup>3</sup>, NYDRA HARVEY-  
COSTELLO<sup>2</sup>, SUSAN HUBBARD<sup>2</sup> AND KENNETH H  
WILLIAMS<sup>2,3</sup>

<sup>1</sup>Lawrence Berkeley Lab

<sup>2</sup>Lawrence Berkeley National Laboratory

<sup>3</sup>Rocky Mountain Biological Laboratory

<sup>4</sup>University of Illinois at Urbana-Champaign

Presenting Author: jwan@lbl.gov

Chemical weathering of silicate rocks can be a sink for atmospheric CO<sub>2</sub>, while weathering of sedimentary rocks can be a source of CO<sub>2</sub> emission. Because sedimentary rocks cover about 2/3 of the Earth's land surface and contain the Earth's largest C inventory as fossil organic carbon (OC<sub>petro</sub>) and inorganic carbonate (IC<sub>petro</sub>), it is important to understand contributions of rock weathering to the terrestrial C cycle. Predictive models on C transfer between rocks and atmosphere are based on the estimated rates of tectonic uplift, weathering, and erosion of exhumated rocks on the Earth's surface over geological time scales. Recent critical zone research has advanced towards greater recognition of subsurface rock weathering. However, quantifying subsurface weathering rates is limited by knowledge of where the most important weathering reactions take place and depth-dependent water fluxes. This research integrates a range of measurements to develop understanding of modern-day rock weathering impacts on C cycling.

This study was conducted in the East River watershed located in the Upper Colorado River Basin where recharge occurs primarily during snowmelt. Along a lower montane hillslope underlain by Mancos Shale, we drilled several deep boreholes and instrumented them with numerous samplers and sensors. We collected 5 years of time- and depth-resolved hydrological and biogeochemical data, including solid, porewater, and pore-gas chemistry, as well as <sup>14</sup>C signatures in solid C, DIC, DOC and CO<sub>2</sub>. We found that (1) modern-day sedimentary rock weathering releases DIC<sub>petro</sub> and DOC<sub>petro</sub> from a narrow region below the soil, which we define as the weathering zone; the annually water table oscillation determines this zone's thickness and the deepest extent of the water table determines the weathering front. (2) The dissolved carbon pool in the subsurface consists primarily of DIC<sub>petro</sub> (~ 85%), which exports through groundwater for eventual CO<sub>2</sub> emission from surface waters (3) The measurements-based annual release rates from shale bedrock weathering are 85 and 15 kg/ha of IC<sub>petro</sub> and OC<sub>petro</sub>, respectively. Our estimated global scale rate of C<sub>petro</sub> release