

# Biological productivity and primary producers recorded in the depositional record of Middle to Late Devonian mudstones from the Horn River Group, Northwest Territories, Canada

MAYA T. LAGRANGE<sup>1</sup>, KAN LI<sup>1</sup>, LONG LI<sup>1</sup>, KURT O. KONHAUSER<sup>1</sup>, BRETTE S. HARRIS<sup>1</sup>, SARA K. BIDDLE<sup>1</sup>, VIKTOR TERLAKY<sup>2</sup> AND MURRAY K. GINGRAS<sup>1</sup>

<sup>1</sup>University of Alberta

<sup>2</sup>Northwest Territories Geological Survey

Presenting Author: mayal@ualberta.ca

The Middle to Late Devonian was characterized by widespread black shale deposition [e.g., 1] and successive marine biotic crises [e.g., 2] attributed to expanded anoxia on marine shelves [e.g., 3]. The cause remains debated and requires constraints from associated marine conditions. This project aims to understand trends in primary producer type and biological productivity of Devonian anoxic pulses. We focus on the Horn River Group (HRG), an Eifelian to Frasnian organic-rich mudstone deposited along the western margin of Laurasia and presently located in the Northwest Territories, Canada. In our study area, this group comprises two units: the Hare Indian Formation and the overlying Canol Formation. The dataset comprises whole-rock  $\delta^{15}\text{N}$  ( $\delta^{15}\text{N}_{\text{bulk}}$ ) and organic  $\delta^{13}\text{C}$  ( $\delta^{13}\text{C}_{\text{org}}$ ) collected from one continuous core.  $\delta^{13}\text{C}_{\text{org}}$  values range from  $-30.4\text{‰}$  to  $-28.2\text{‰}$  and show a peak in the Hare Indian Formation with a second spike in the Canol Formation.  $\delta^{15}\text{N}_{\text{bulk}}$  values are narrow in range ( $-3.7\text{‰}$  to  $+1.9\text{‰}$ ), without observable trends in the HRG. Nitrogen isotopic ratios near  $0\text{‰}$  are characteristic of  $\text{N}_2$ -fixing (diazotrophic) primary producers and thus indicate that diazotrophs were dominant throughout deposition of the HRG, despite fluctuations in marine paleoredox. Trends in  $\delta^{13}\text{C}_{\text{org}}$  throughout the Canol Formation indicate variations in biological productivity, suggesting that locally the growth rate of primary producers varied alongside changes in paleoredox. Our results can be compared to age-equivalent black shales to understand regional and global ecological patterns that contributed to widespread organic carbon burial.

[1] Klemme, H.D., Ulmishek, G.F., 1991. Effective Petroleum Source Rocks of the World: Stratigraphic Distribution and Controlling Depositional Factors. *Am. Assoc. Pet. Geol. Bull.* 75, 1809–1851.

[2] Walliser, O.H., 1996. Global Events in the Devonian and Carboniferous, in Walliser, O.H. (Ed.), *Global Events and Event Stratigraphy in the Phanerozoic*. Springer Berlin Heidelberg, pp. 225–250.

[3] Lu, M., Lu, Y., Ikejiri, T., Sun, D., Carroll, R., Blair, E.H., Algeo, T.J., Sun, Y., 2021. Periodic oceanic euxinia and terrestrial fluxes linked to astronomical forcing during the Late Devonian Frasnian–Famennian mass extinction. *Earth Planet. Sci. Lett.* 562, 116839.