The coming and going of banded iron formation

MS. RUIMIN WANG, PHD STUDENT 1 AND BING SHEN^2

¹Peking University ²School of Earth and Space Sciences, Peking University Presenting Author: scorpionwrm@pku.edu.cn

The banded iron formation (BIF) is the signature of Archean and Paleoproterozoic deposits, disappeared in 1.8 billion years ago (Ga) marked a sharp change of environments. It has long been held that the disappearance of BIF might be related to the seawater depletion of Fe²⁺(ferrous iron), probably due to the ocean oxidation or the development of oceanic euxinia. However, more and more geochemical and sedimentological evidence indicates that the Proterozoic ocean was predominant ferruginous (Fe2+ enriched) with the widespread sulfidic continental margins, and non-BIF ironstone deposits are common after 1.8 Ga, questioning the Fe-depletion interpretation for the disappearance of BIF. Here, we show the genetic relationship between Fe, Si, and P mineral phases, suggesting the coupled Fe-Si-P cycles in BIF formation. The Fe redox cycle and Si cycle are coupled by the scavenging of seawater Si by FeOOH precipitation. We suggest active Fe redox cycle after 1.8 Ga might have enhanced the efficiency of silica precipitation from seawater, lowering the seawater silica concentration. As such, BIF precipitation was prohibited by the insufficient supply of silica from seawater. In contrast, the non-BIF ironstone deposition was warranted by iron supply from ferruginous seawater. This model can also explain the reoccurrence of BIF in the Neoproterozoic Snowball Earth (~0.7 Ga), during which inactive Fe-redox cycle allowed the increase of seawater silica concentration in the ice-covered ocean.