

Kinetic biogeochemical modeling of a high-pressure reactor simulating H₂ geological storages in deep aquifers.

JEAN MURA¹, MAGALI RANCHOPEYRUSE^{1,2,3},
MARION GUIGNARD³, MARION DUCOUSSO^{1,2}, MARIE
LARREGIEU³, PASCALE SÉNÉCHAL⁴, MARIE-PIERRE
ISAURE³, ISABELLE LE HÉCHO^{2,3}, PETER MOONEN⁴,
GUILHEM HOAREAU⁵, MATEUS BURITI¹, MARIE
POULAIN¹, ANÉLIA PETIT⁶, PIERRE CHIQUET^{2,7},
GUILHEM CAUMETTE^{2,8}, PIERRE CÉZAC^{1,2} AND
ANTHONY RANCHOPEYRUSE^{2,3}

¹Université de Pau et des Pays de l'Adour, E2S UPPA, LaTEP

²Joint Laboratory SEnGA, UPPA-E2S-Teréga

³Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, IPREM

⁴Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, DMEX

⁵Université de Pau et des Pays de l'Adour, E2S UPPA, CNRS, TOTAL, LFCR

⁶STORENGY – Geosciences Department

⁷Teréga – Geosciences Department

⁸Teréga – Environment Department

Presenting Author: jean.mura@univ-pau.fr

If dihydrogen (H₂) is to play a key role in tomorrow's energy mix, massive storage quantities will be required, which can only be achieved by using underground gas storages (UGS), such as deep aquifers.

The main challenge expected with H₂ storage in aquifers is its consumption by microorganisms. Besides consuming the stored energy, these reactions could harm the storage by pressure modification, souring, and modified porosity and permeability via precipitation and dissolution of minerals, as well as bio-clogging. Both recent conceptual modeling and experimental studies underline the need for more experimental data to fit models and upscale them to the aquifer scale.

This work aimed to assess the impact of injecting 2% H₂ over three months into a three-phase high-pressure bioreactor [1,2] simulating the conditions of a deep aquifer with high sulfate concentration (9 mmol/kgw). The built-in capabilities of the PHREEQC software were used to model the liquid-gas and liquid-solid equilibrium, acido-basic, and complexation reactions in a 0D model. New kinetic laws were proposed and implemented into PHREEQC based on microbial taxonomic diversity analyses to model the intricate biotic phenomena observed. This model allows us to represent the growth and decay of the microbial population, respiration rates, and multiple inhibitions.

During this experiment, sulfate reduction occurred until total sulfate consumption, followed by methanogenesis. Formate and acetate production also happened at a slower rate. These reactions consumed 76% of the H₂ quantity injected. However, while H₂ remained in sufficient quantity, the microbial activity

stopped after the total consumption of all electron acceptors and alkalization (pH 9.5). We hypothesize competition for CO₂/carbonates between microorganisms and geochemical processes (*i.e.*, mineral precipitation). This suggested that H₂ could be stable in this aquifer after an initial loss. Although sulfate reduction induced notable sulfide production, all the sulfide produced was attenuated, probably through the precipitation of pyrite. The developed model represents a preliminary step toward a comprehensive description of H₂ fate in the underground, allowing for an upscaled reservoir model in a second step.

[1] Haddad et al. 2022, Energy Environ. Sci. 15, 3400.

[2] Mura et al. 2024, International Journal of Hydrogen Energy. 63, 330.