

Effect of deformation and metamorphism on metal redistribution at the Sulitjelma volcanogenic massive sulfide deposits

ESZTER SENDULA¹, CARLY FABER¹, LEIF-ERIK R. PEDERSEN², EDUARDO TEIXEIRA MANSUR³, TATSUO NOZAKI⁴, HIRONAO MATSUMOTO⁴ AND SABINA STRMIC PALINKAS¹

¹UiT – The Arctic University of Norway

²Department of Earth Science, Center for Deep Sea Research, University of Bergen, Bergen, Norway

³The Geological Survey of Norway (NGU)

⁴Japan Agency for Marine-Earth Science and Technology (JAMSTEC)

Presenting Author: eszter.sendula@uit.no

The Sulitjelma historical mining district in northern Norway shows classic examples of Cu-(Zn) volcanogenic massive sulfide (VMS) deposits, containing more than 20 known ore bodies. The Cu-(Zn) mineralization formed in a marginal back-arc basin setting during Ordovician westward subduction at the Laurentia margin. The basin experienced closure, with ophiolite obduction, as a result of the Scandian phase of the Caledonian Orogeny at 430-420 Ma¹. The observed greenschist to amphibolite facies metamorphism is related mostly to Scandian phase deformation and subsequent exhumation.

Deformation and metamorphism can play a significant role in modification of primary deposits and (re)mobilization of metals and elements in sulfide minerals, however the role of crystal plastic deformation in mobilization is not often investigated and therefore poorly understood. Dubosq et al.² showed that several trace elements are introduced or remobilized into deformation-induced substructures within pyrite during deformation. Metamorphic reactions are commonly accompanied by generation of fluids of various compositions which can influence the nature and extent of deformation mechanisms and element remobilization. Our work investigates the interplay between deformation, element remobilization and fluids during greenschist- to amphibolite-facies metamorphism on the VMS deposits at Sulitjelma using both geochemical and microstructural mapping methods to characterize multiphase mobilization of sulfides.

Samples were collected from the Ny-Sulitjelma, Hankabakken, Giken, Jakobsbakken and Sagmo ore deposits and were studied by optical light microscopy, SEM, EBSD, Raman spectroscopy, trace element mapping by LA-ICP-MS, and fluid inclusion microthermometry. We will present preliminary results combining EBSD and trace element data on pyrite grains, showing a complex history of the ores in the area. Numerous fluid inclusion assemblages (FIA) were found with variable compositions (salinity ranging from 0.9±0.3 wt% NaCl_{equiv} to 44.3±1 wt% NaCl_{equiv}; presence of different volatiles including CO₂, N₂ and CH₄) and homogenization temperatures (between

138±7°C and 363±5°C) reflecting the evolution of metamorphic fluids that contributed to remobilization in metals during the metamorphic phase of the Sulitjelma deposits.

¹Roberts, D., and Sturt, B.A. (1980). *J. Geol. Soc.*, London, 137, p. 241-250.

²Dubosq, R., Lawley, C.J.M., Rogowitz, A., Schneider, D.A., and Jackson, S. (2018). *Lithos*, 310-311, p. 86-104.