

# Constraints on Weathering Spikes by Stable Tungsten Isotope during the Past 300,000 Years

SHU LIU<sup>1</sup>, RONG HU<sup>1</sup>, TAO LI<sup>2</sup>, RUIYU YANG<sup>3</sup> AND  
GAOJUN LI<sup>1</sup>

<sup>1</sup>Nanjing University

<sup>2</sup>Nanjing Institute of Geology and Palaeontology, Chinese Academy of Sciences

<sup>3</sup>University of Cologne

Presenting Author: lkr2000ls@163.com

How the continental weathering fluxes have responded to the Quaternary glacial-interglacial (G-IG) climate cycles is debated, which is critical to understand the effects of climate change on weathering. With a residence time of ~4,000 years and riverine input as the dominant source<sup>[1]</sup>, tungsten(W) and its stable isotope composition ( $\delta^{186/184}\text{W}$ ) in the seawater could hopefully document the variation of weathering flux in the glacial-interglacial timescale. Here, we present a stable W isotope record of ocean sediments from ODP Site 834A at Lau Basin on a high temporal resolution during the past 300,000 years to constrain the weathering flux changes. Our record shows that while the  $\delta^{186/184}\text{W}$  of seawater remained stable overall in the studied period, rapid fluctuations up to 0.1‰ occurred during major G-IG transitions. Based on a simplified mass balance model, we propose that the documented oceanic  $\delta^{186/184}\text{W}$  shifts were driven by the “weathering spikes” following the glacial terminations, corresponding to 5~10 fold increase in weathering fluxes. This suggests that W isotope has the potential to investigate the relationship between continental weathering and climate change.

[1]R. Yang et al. (2022), *Geochimica et Cosmochimica Acta* 322, 227-243.