

# Stable Sr isotopic variability in Godavari River basin, India: implications for surface processes and reservoir mixing

SOURAV GANGULY<sup>1</sup>, VENKATA SAILAJA PAPPALA<sup>2</sup>  
AND RAMANANDA CHAKRABARTI<sup>1</sup>

<sup>1</sup>Indian Institute of Science

<sup>2</sup>North Carolina State University

Presenting Author: sganguly.work@gmail.com

Understanding the chemistry of dissolved constituents of rivers is critical to constrain a variety of Earth surface processes that play major role in modulating global elemental cycles. Stable Sr isotopes have emerged as a new tool to better understand low-temperature surface processes. Here we present new stable Sr isotope data from a large silicate-draining river basin of southern India and discuss possible implications. Water samples were collected from the Godavari River near Rajahmundry (~ 65 Km from the coast), Andhra Pradesh during monsoon (wet) and non-monsoon (dry) months of 2015-16. The dissolved cation concentrations were measured using an ICP-MS at the Centre for Earth Sciences, Indian Institute of Science and the external precision was better than  $\pm 5\%$ . The radiogenic ( $^{87}\text{Sr}/^{86}\text{Sr}$ ) and stable Sr ( $\delta^{88/86}\text{Sr}$ ) isotopic measurements were carried out using established protocols<sup>1,2</sup> using a TIMS at Centre for Earth Sciences, Indian Institute of Science. External precisions (2SD) for radiogenic and stable Sr isotope measurements were better than  $\pm 15$  ppm and  $\pm 0.024\%$  based on multiple measurements of NIST SRM-987 standard. The dissolved Sr content in the river water samples varied between 0.55 - 1.53  $\mu\text{mol/L}$ . Radiogenic  $^{87}\text{Sr}/^{86}\text{Sr}$  ranged between 0.714249 - 0.719358 while the  $\delta^{88/86}\text{Sr}$  varied between 0.174 - 0.640 ‰. The average  $\delta^{88/86}\text{Sr}$  of Godavari river is estimated at  $0.327 \pm 0.226\%$  (2SD, N = 18) which matches with global average  $\delta^{88/86}\text{Sr}$  of world rivers. When compared with the  $^{87}\text{Sr}/^{86}\text{Sr}$  data the samples essentially overlapped with the bulk silicate earth estimates underscoring the dominance of congruent weathering of surrounding lithologies. Although a distinct seasonal trend was lacking, the heaviest  $\delta^{88/86}\text{Sr}$  values were observed mostly in the dry seasons which suggests mixing with a fractionated reservoir such as the local groundwater, specifically when contributions from seawater was minimal.

[1] Banerjee et al. (2016) Chemical Geology 440,124-138.

[2] Ganguly and Chakrabarti (2022) Journal of Analytical Atomic Spectrometry 37(10),1961-1971.