

Assessment of CO₂ consumption from basalt amendments to soils.

LOUIS A DERRY^{1,2}, OLIVER CHADWICK³ AND
STEPHEN PORDER⁴

¹Institut de Physique du Globe de Paris

²Cornell University

³University of California Santa Barbara

⁴Brown University

Presenting Author: derry@cornell.edu

Concern over the impacts of anthropogenic greenhouse gas emissions has led to proposals to offset CO₂ emissions by mechanisms that could increase carbon storage in soils. Estimates of the potential efficacy of various strategies vary, but an issue common to all of them is how to verify the magnitude and stability of the results of interventions. Soil sinks are “out of sight” which contrasts with strategies like reforestation. Below-ground carbon sinks, whether they be organic or inorganic, pose substantially more challenging issues for quantification and monitoring.

ERW seeks to apply Ca, Mg-rich silicate rocks to enhance the consumption of CO₂ by weathering reactions. Measurement of base cation losses from soils is used for quantifying CO₂ uptake. Assessing CO₂ uptake this way requires assessment of small differences between heterogeneous end members. Geochemical tracers can be used to estimate basalt input assuming that the endmembers are distinct. To compensate for open system behavior normalization to an “immobile” element is necessary. The limitation is typically the highly heterogeneous nature of soils. Data from these settings often have high covariances. We reanalyzed published data on amended soils using Monte Carlo uncertainty analysis. We find that in a number of cases the Ca and Mg differences in pre- and post-amendment soils are not significantly different from zero (1 s.e.). High soil chemistry variances makes quantification of small differences difficult.

Techniques for estimating relevant sample size and power for noisy data sets and modest effect sizes are well developed in other fields and can be appropriately applied to ERW problems. A simplified example using Lehr’s approximate rule for a two-sided test with s.d. for the pre- and post- data sets = 0.1 and the effect size (net change) = 0.03 yields a sample size $n = 178$ to obtain a sample power of 0.8 at the 95% CI, an optimistic estimate. Appropriate experimental design for ERW will require substantial sampling effort and rigorous *a priori* statistical assessment. The ocean $\Delta\text{CO}_2/\Delta\text{ALK}$ ratio used to estimate CO₂ uptake is important and unlikely to be as high as most studies have assumed.